

Physical Chemistry Laboratory I

Experiment 3

Effect of Ionic Strength on the Solubility of CaSO₄


(Revised, 01/13/03)

It is generally assumed that solutions of strong electrolytes are completely dissociated and that the deviations from ideal behavior (in freezing point depression or conductivity) of the electrolyte solutions result from interionic attractions of the ions. However, the early literature is replete with examples of partially dissociated species, or ion pairs for polyvalent electrolytes, even in moderately dilute solutions. There are many recent articles on the dissociation constants of ionic compounds in aqueous solutions. Ion pairs are important for polyvalent electrolytes in water and in media of low dielectric constant (water/organic solvent mixtures or water at high temperatures).

The standard treatment of solubility and solubility product constants in introductory chemistry courses is generally doubly simplified: 1st by the assumption that the activity coefficients are 1, and 2nd by the assumption that the ionic salts are completely dissociated in solution. Both of these assumptions are reasonable for very slightly soluble 1/1 electrolytes (such as AgCl), but neither is correct for somewhat more soluble polyvalent electrolytes. Some recent texts for quantitative analysis include significant discussions of activity coefficients and introduce the concept of ion pairs for polyelectrolytes. [1]

The solubility of CaSO₄ and its variation with concentration of added electrolyte illustrate the effects of activity coefficients and ion pairs. There have been many articles in Journal of Chemical Education (J. Chem. Ed.) that discuss the solubility and the use of solubility data to calculate K_{SP}(CaSO₄). [2 - 7] Ion pairing is not as well known as it should be. Although we will be concerned only with the interactions of Ca²⁺ with SO₄²⁻ in this experiment, non-covalent bonding of Ca²⁺ is frequently observed with other species.

The dissociation reaction that is often used to describe the solubility of CaSO₄ in water is the following, which assumes that calcium sulfate is completely dissociated in aqueous solution:



From this equation, one may write the appropriate equilibrium expression, including activity coefficient effects:

$$K_{\text{SP}}^0 = a\{\text{Ca}^{2+}\} \cdot a\{\text{SO}_4^{2-}\} = [\text{Ca}^{2+}] [\text{SO}_4^{2-}] \gamma_{\pm}^2 \quad (2a)$$

$$K_{\text{SP}}^0 = K_{\text{SP}}\{\text{Apparent}\} \gamma_{\pm}^2 = S^2 \gamma_{\pm}^2 \quad (2b)$$

In these equations, $a\{X\}$ = activity of X; the concentration of each species, $[X_i]$, is given in mol $\{X_i\}/L$; γ_{\pm} refers to the mean (geometric average) activity coefficient for the two ions ($\gamma_{\pm}^2 = \gamma\{\text{Ca}^{2+}\} \cdot \gamma\{\text{SO}_4^{2-}\}$); K_{SP}^0 is the "true" solubility product constant for CaSO₄, i. e., the value at infinite dilution or "zero concentration" of ions; and $K_{\text{SP}}\{\text{Apparent}\}$ is the product of the ionic concentrations, $[\text{Ca}^{2+}] \cdot [\text{SO}_4^{2-}]$. With this model, there is no other source of either ion when pure calcium sulfate is dissolved in water; therefore, $[\text{Ca}^{2+}] = [\text{SO}_4^{2-}] = S(\text{CaSO}_4)$.

S{CaSO₄} in pure water at 25.0 °C.

The determination of the solubility of a dense solid like $\text{CaSO}_4(\text{s})$ is not particularly difficult at room temperature. Add some $\text{CaSO}_4(\text{s})$ to a 250 mL Erlenmeyer flask add ~ 200 mL of distilled H_2O , immerse in the constant temperature bath at 25.0°C for ~ 15 minutes, with stirring: magnetic stirrer and stirring bar. Use the lead weight around the flask to weigh it down. Neither the weight of calcium sulfate nor the volume of water is critical as long as solid is present at the end of the experiment. However, record the weight of $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ that you use. Calculate the amount of $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ needed to dissolve in 200 mL of water and use $\sim 100\%$ excess.

Let the solution settle for ~ 15 minutes in the constant temperature bath and then use a volumetric pipet to take a 25.00 mL aliquot (sample) of the solution. (Use the automatic pipetter.) Standard quantitative procedures require that you rinse the pipet with the solution being analyzed before taking the sample. Don't put the pipet on the bottom and remove solid CaSO_4 .

Transfer the aliquot to a beaker, add ~ 50 mL of water (graduated cylinder OK), a pinch of the solid hydroxynaphthol blue (HNB) indicator, and adjust the pH of the solution to ~ 11.0 with ~ 5 mL of $\text{NH}_3/\text{NH}_4\text{Cl}$ buffer (graduated cylinder OK). Use pH meter. (See the [*Appendix*](#) for general information about the pH meter. Operating instructions are provided.) If necessary, increase the pH by adding a few drops of 1 M NaOH. Titrate the solution with standard EDTA solution to the appearance of a pure blue. A carboy of standardized $\text{Na}_2\text{H}_2\text{EDTA}$ is available. Do the titration in a beaker, so that you can keep the pH electrode in the solution during the titration. Be careful that the stirring bar does not hit the pH electrode. If the pH drops below 10 during the titration, add a few mL of the buffer to return the pH to ~ 11 .

Duplicate samples should be analyzed. Use both values in your analysis of the data (not the average). $S(\text{CaSO}_4, 25^\circ\text{C}) \approx 0.015\text{ M}$. If you obtain significantly different values, check calculations and/or repeat the titration. The solubility of CaSO_4 does not vary much around room temperature. This analysis gives the total amount of calcium in solution in all forms: $\text{Ca}^{2+}(\text{aq})$ and $\text{CaSO}_4(\text{aq})$, if it is present. Experiments reported in the literature have sometimes involved very long equilibration times. However, long times do not seem to be necessary for experiments at room temperature.

EDTA titrations are discussed in Harris, *Quantitative Chemical Analysis*, or other recent quantitative analysis texts. One mole of EDTA reacts with one mole of Ca^{2+} . The end point of the titration is the appearance of a pure blue in a solution that was initially a wine red. This end point is not as easy to see as the phenolphthalein end point for the titration of HCl with NaOH in Exp. 1. However, accurate values for $[\text{Ca}^{2+}]$ can be achieved using this standard procedure.

One way to recognize the end point color is to take ~ 50 mL of distilled water, bring to pH ~ 11 , add a few drops of EDTA solution (to complex any cations that might be present as impurities), and add a pinch of the indicator and observe the color. Keep this solution (or one of the titrated solutions) as a reference for the blue color.

One can use the Debye-Huckel-Guggenheim (DHG) equation to calculate activity coefficients for solutions of known concentration (ionic strength). (Noggle, Chap. 8; Tinoco, Jr., Sauer, & Wang, Chap. 4)

$$\log \gamma_j = -\left(\frac{0.509 Z_j^2 \sqrt{I}}{1 + \sqrt{I}}\right) \quad \text{or} \quad \log \gamma_{\pm} = -\left(\frac{0.509 |Z_+ Z_-| \sqrt{I}}{1 + \sqrt{I}}\right) \quad (3)$$

In this equation, the ionic strength, I or μ , is given by the following equation.

$$\mu = I = \frac{\sum_j m_j Z_j^2}{2} \quad (4)$$

The ionic strength includes all ions in the solution; Z_i is the charge of an ion of concentration, m ; and the constant, 0.509, applies to aqueous solutions at 25.0 °C. This equation gives an activity coefficient of 1 for an ion pair, because $Z = 0$. The DHG equation actually uses molality, m , in calculating the ionic strength, I . For dilute aqueous solutions like this one, $m \approx M$ and no significant error is introduced by using molarity in DHG equation, < 1%.

If one assumes that there are no ion pairs in solution, the $S\{\text{CaSO}_4\} = [\text{Ca}^{2+}] = [\text{SO}_4^{2-}]$ and Eq. (2b) applies. With activity coefficients calculated with the DHG equation, Eq. (3), one can calculate K_{SP}° from the solubility of CaSO_4 in pure water, if there are no ion pairs in solution.

Calculate K_{SP}° from these solubility data and compare with literature values. Report these results in a table. Compare this value with $K_{\text{SP}}\{\text{Apparent}\} = [\text{Ca}^{2+}][\text{SO}_4^{2-}] = S^2$.

Effect of ionic strength on $S\{\text{CaSO}_4\}$ at 25.0 °C

In separate 250 mL volumetric flasks prepare five solutions of an inert electrolyte, NaCl, of accurately known concentrations. The concentrations should be in the range ~ 0.05 M to ~ 0.25 M. The exact values are not critical but each solution should be known with an accuracy of ~ one part in 1000. A volume of 250 mL of each solution should be sufficient. Use the analytical balance and weigh NaCl to ± 0.0001 g.

Repeat the previous procedure to determine the solubility of CaSO_4 in water with each of these salt solutions. Add ~ 200 mL of each solution to excess $\text{CaSO}_4(\text{s})$ in a 250 mL Erlenmeyer flask. Stir the solution (~ 15 minutes) and allow it to settle in the constant temperature bath (~ 15 minutes) as before. These concentrations of added inert electrolyte are such that $S(\text{CaSO}_4)$ will be significantly larger than the value in pure water. Duplicate titrations of each solution should be done. Use both values in the analysis of your data.

You should record your data in a table as $C\{\text{NaCl}, M\}$ and $S\{\text{CaSO}_4, M\}$. The values for the concentrations of NaCl should be given to four decimal places and the solubility of CaSO_4 should be given to four significant figures (five decimal places). Plot these results as $S\{\text{CaSO}_4, M\}$ vs. $C\{\text{NaCl}, M\}$. The solubility of CaSO_4 should increase monotonically with increasing $M\{\text{salt}\}$. Include your values for $S\{\text{CaSO}_4\}$ with no added inert electrolyte: $C\{\text{NaCl}, M\} = 0$.

This solution has the smallest value of ionic strength (I or μ) that you can obtain. However, the activity coefficients are not unity for this solution (as you calculated above).

In principle, you can determine the molality of each solution because you know the concentration of the added salt and that value is larger than the concentration of dissolved calcium sulfate. An equation relating the activity coefficients in molal units, γ , and in molar units, y , is known, $y_{\pm} = \gamma_{\pm}(m/M)d_o$. [9] In this equation, d_o is the density of the solvent. From data in the CRC Handbook of Chemistry and Physics, the ratio of molality to molarity, $m/M \leq 1.01$ for $M \leq 0.2$ mol/L (mol/dm³) for several alkali metal salts. Consequently, you may use the DHG equation to calculate activity coefficients using the molarities that you determine from your titration and introduce no significant error in the analysis.

Following Ramette [2], one can analyze the data by neglecting ion pairs (as was done in the previous section above) and calculate $K_{SP}\{\text{apparent}\}$ from the solubility data and extrapolate to $I = 0$ to obtain K_{SP}^0 . Extrapolations are done as linear functions whenever possible. If one takes the negative logarithms of both sides of Equation (2b) and rearranges, one obtains

$$-\log\{S^2\} = pK_{SP}\{\text{Apparent}\} = pK_{SP}^0 + 2*\log\{\gamma_{\pm}\} \quad (5)$$

Use Equation (3) to calculate $\log\{\gamma_{\pm}\}$ from the ionic strength at each concentration. Remember to include all ions, Ca^{2+} , SO_4^{2-} , Na^+ , and Cl^- in calculating I (or μ).

Plot $pK_{SP}\{\text{Apparent}\}$, or $-\log\{S^2\}$, vs $\log\{\gamma_{\pm}\}$ and extrapolate to $\log\{\gamma_{\pm}\} = 0$ to get the intercept, pK_{SP}^0 . {In the somewhat confusing and perhaps unnecessary way, $pK_{SP} = -\log K_{SP}$, by analogy with pH.} There are other functions that you can plot; however, this plot shows how the apparent pK_{SP} varies with ionic strength. Use a spreadsheet (Excel, etc), Mathcad, or Maple for the plot (which should be a part of your report). Compare this value with pK_{SP}^0 values that you obtained from the literature (a quantitative analysis text or a primary reference) and with the value that you obtained from the solubility of calcium sulfate in pure water. Compare the slope of your experimental curve with the expected value from Equation (5).

If significant concentrations of ion pairs of $CaSO_4$ exist in aqueous solutions, then equation (2a) is still correct, but the approximation $S = [Ca^{2+}] = [SO_4^{2-}]$ is not correct and Equation (2b) is incorrect. The solubility of $CaSO_4$ is given by a two-term equation,

$$S = [Ca^{2+}] + [CaSO_4]. \quad (6)$$

In equation (6), $[Ca^{2+}]$ = the molarity of Ca^{2+} ions in solution and $[CaSO_4]$ = the concentration of the ion pair, M .

Equation (2a) is still correct and the stoichiometric relationship $[Ca^{2+}] = [SO_4^{2-}]$ is still correct. Consequently,

$$[Ca^{2+}] = \sqrt{(K_{SP}^0/\gamma_{\pm}^2)} = \sqrt{(K_{SP}^0)}/\gamma_{\pm} \quad (7)$$

The additional reaction and equilibrium expression involve the dissociation of the ion pair,



The dissociation equilibrium constant for this chemical reaction is given in (9)

$$K_D = \frac{[\text{Ca}^{2+}, \text{aq}][\text{SO}_4^{2-}, \text{aq}]\gamma_{\pm}^2}{[\text{CaSO}_4, \text{aq}]} \quad (9)$$

Solving Equation (9) for $[\text{CaSO}_4, \text{aq}]$, one obtains the following expression,

$$[\text{CaSO}_4, \text{aq}] = \frac{[\text{Ca}^{2+}, \text{aq}][\text{SO}_4^{2-}, \text{aq}]\gamma_{\pm}^2}{K_D} = \frac{K_{SP}^o}{K_D} \quad (10)$$

Substituting this relationship and Equation (7) into Equation (6), one obtains (finally)

$$S(\text{CaSO}_4, M) = \frac{\sqrt{K_{SP}^o}}{\gamma_{\pm}} + \frac{K_{SP}^o}{K_D} \quad (11)$$

Plot your data as S vs $1/\gamma_{\pm}$ to get values for the slope, $\sqrt{K_{SP}^o}$, and the intercept, K_{SP}^o/K_D . A plot of your data should give a reasonably straight line with a positive intercept. The value that you obtain for K_{SP}^o by this method may not be the same as the value you obtained using Eq. (5).

There is, however, one problem with these values for K_{SP}^o and K_D : they are only first approximations to the correct values. The activity coefficients, γ_{\pm} , were calculated as if $[\text{Ca}^{2+}] = [\text{SO}_4^{2-}] = S$. Although $[\text{Ca}^{2+}] = [\text{SO}_4^{2-}]$ is still correct, the analysis must be done again by calculating the activity coefficients, γ_{\pm} , using the better estimate that

$$[\text{Ca}^{2+}] = [\text{SO}_4^{2-}] = S - \frac{K_{SP}^o}{K_D} \quad (12)$$

Replot and reanalyze the data as S vs $1/\gamma_{\pm}$ with the corrected activity coefficients to obtain more reliable values for the intercept and slope. Compare your results with literature values.

The following data were reported in 1918 on the solubility of CaSO_4 in solutions of KNO_3 at 25.0 °C.

KNO_3, M	$S(\text{CaSO}_4), M$
0	0.01530
0.02755	0.01805
0.05268	0.02008
0.10307	0.02120

Are these data consistent with your solubility data? To the approximation given by the activity coefficient equations, the chemical nature of the ions should have no effect at the solubility. Analyze these data to obtain K_{SP}^o and K_D .

References

1. D. C. Harris, Quantitative Chemical Analysis, 5th Ed., 1999, Chap. 8
2. R. W. Ramette, *J. Chem. Ed.* 33, 610 (1956).
3. L. Meites, J. S. F. Pode, and H. C. Thomas, *J. Chem. Ed.* 43, 667 (1966).
4. A. K. Sawyer, *J. Chem. Ed.* 60, 416 (1983).
5. R. B. Martin, *J. Chem. Ed.* 63, 471 (1986).
6. D. Masterman, *J. Chem. Ed.* 64, 409 (1987).
7. S. O. Russo and G. I. H. Hanania, *J. Chem. Ed.* 66, 148 (1989)
8. J. H. Carpenter, *J. Chem. Ed.* 66, 184 (1989).
9. R. A. Robinson and R. H. Stokes, Electrolyte Solutions, Butterworths, 1955, p. 30.
10. R. G. Ainsworth, *J. Chem. Soc. Farad*, 69, 1028 (1973).

Appendix

GENERAL pH METER INSTRUCTIONS

PLEASE TREAT THIS EXPENSIVE EQUIPMENT WITH CARE

- The measuring electrode is fragile: do not stir with it; do not allow a stirring bar to hit it; do not jam it into bottom of any container.
- Unscrew the plastic storage container to remove it from the electrode, leaving the cap on the electrode. Do not pull the container or container cap off of the electrode.
- Always rinse the electrode with distilled water when moving it to a different solution or buffer. Rinse, but do not wipe or blot it.
- The white ring at the top of electrode must remain in its closed position.
- The electrode must **NEVER** dry out: it must always be either in the solution being measured, or in the filled storage container!
- At the end of your experiment session, as part of your clean up, empty the storage container, rinse it out with distilled water, refill it (to about $\frac{3}{4}$ -full) with the KCl storage solution provided, and then return the electrode into the container, standing upright.
- To properly standardize the meter, refer to instructions in the **Operation Manual** at the station for the experiment.