

# Chapter F01

## Matrix Factorizations

### Contents

<b>1</b>	<b>Scope of the Chapter</b>	<b>2</b>
<b>2</b>	<b>Background to the Problems</b>	<b>2</b>
2.1	Matrix Inversion . . . . .	2
2.2	Matrix Factorizations . . . . .	3
2.3	Matrix Arithmetic and Manipulation . . . . .	3
<b>3</b>	<b>Recommendations on Choice and Use of Available Routines</b>	<b>4</b>
3.1	Matrix Inversion . . . . .	4
3.2	Matrix Factorizations . . . . .	5
3.3	Matrix Arithmetic and Manipulation . . . . .	5
<b>4</b>	<b>Decision Trees</b>	<b>5</b>
<b>5</b>	<b>Index</b>	<b>8</b>
<b>6</b>	<b>Routines Withdrawn or Scheduled for Withdrawal</b>	<b>9</b>
<b>7</b>	<b>References</b>	<b>9</b>

# 1 Scope of the Chapter

This chapter provides facilities for three types of problem:

- (i) Matrix Inversion
- (ii) Matrix Factorizations
- (iii) Matrix Arithmetic and Manipulation

These problems are discussed separately in Section 2.1, Section 2.2 and Section 2.3.

## 2 Background to the Problems

### 2.1 Matrix Inversion

- (i) Non-singular square matrices of order  $n$ .

If  $A$ , a square matrix of order  $n$ , is non-singular (has rank  $n$ ), then its inverse  $X$  exists and satisfies the equations  $AX = XA = I$  (the identity or unit matrix).

It is worth noting that if  $AX - I = R$ , so that  $R$  is the ‘residual’ matrix, then a bound on the relative error is given by  $\|R\|$ , i.e.,

$$\frac{\|X - A^{-1}\|}{\|A^{-1}\|} \leq \|R\|.$$

- (ii) General real rectangular matrices.

A real matrix  $A$  has no inverse if it is square ( $n$  by  $n$ ) and singular (has rank  $< n$ ), or if it is of shape ( $m$  by  $n$ ) with  $m \neq n$ , but there is a **Generalized** or **Pseudo Inverse**  $Z$  which satisfies the equations

$$AZA = A, \quad ZAZ = Z, \quad (AZ)^T = AZ, \quad (ZA)^T = ZA$$

(which of course are also satisfied by the inverse  $X$  of  $A$  if  $A$  is square and non-singular).

- (a) if  $m \geq n$  and  $\text{rank}(A) = n$  then  $A$  can be factorized using a **QR factorization**, given by

$$A = Q \begin{pmatrix} R \\ 0 \end{pmatrix},$$

where  $Q$  is an  $m$  by  $m$  orthogonal matrix and  $R$  is an  $n$  by  $n$ , non-singular, upper triangular matrix. The pseudo-inverse of  $A$  is then given by

$$Z = R^{-1} \tilde{Q}^T,$$

where  $\tilde{Q}$  consists of the first  $n$  columns of  $Q$ .

- (b) if  $m \leq n$  and  $\text{rank}(A) = m$  then  $A$  can be factorized using an **RQ factorization**, given by

$$A = (R \ 0) P^T$$

where  $P$  is an  $n$  by  $n$  orthogonal matrix and  $R$  is an  $m$  by  $m$ , non-singular, upper triangular matrix. The pseudo-inverse of  $A$  is then given by

$$Z = \tilde{P} R^{-1},$$

where  $\tilde{P}$  consists of the first  $m$  columns of  $P$ .

- (c) if  $m \geq n$  and  $\text{rank}(A) = r \leq n$  then  $A$  can be factorized using a **QR factorization**, with column interchanges, as

$$A = Q \begin{pmatrix} R \\ 0 \end{pmatrix} P^T,$$

where  $Q$  is an  $m$  by  $m$  orthogonal matrix,  $R$  is an  $r$  by  $n$  upper trapezoidal matrix and  $P$  is an  $n$  by  $n$  permutation matrix. The pseudo inverse of  $A$  is then given by

$$Z = P R^T (R R^T)^{-1} \tilde{Q}^T,$$

where  $\tilde{Q}$  consists of the first  $r$  columns of  $Q$ .

- (d) if  $\text{rank}(A) = r \leq k = \min(m, n)$ , then  $A$  can be factorized as the **singular value decomposition**

$$A = QDP^T,$$

where  $Q$  is an  $m$  by  $m$  orthogonal matrix,  $P$  is an  $n$  by  $n$  orthogonal matrix and  $D$  is an  $m$  by  $n$  diagonal matrix with non-negative diagonal elements. The first  $k$  columns of  $Q$  and  $P$  are the **left-** and **right-hand singular vectors** of  $A$  respectively and the  $k$  diagonal elements of  $D$  are the **singular values** of  $A$ .  $D$  may be chosen so that

$$d_1 \geq d_2 \geq \dots \geq d_k \geq 0$$

and in this case if  $\text{rank}(A) = r$  then

$$d_1 \geq d_2 \geq \dots \geq d_r > 0, \quad d_{r+1} = \dots = d_k = 0.$$

If  $\tilde{Q}$  and  $\tilde{P}$  consist of the first  $r$  columns of  $Q$  and  $P$  respectively and  $\Sigma$  is an  $r$  by  $r$  diagonal matrix with diagonal elements  $d_1, d_2, \dots, d_r$  then  $A$  is given by

$$A = \tilde{Q}\Sigma\tilde{P}^T$$

and the pseudo inverse of  $A$  is given by

$$Z = \tilde{P}\Sigma^{-1}\tilde{Q}^T.$$

Notice that

$$A^T A = P(D^T D)P^T$$

which is the classical eigenvalue (spectral) factorization of  $A^T A$ .

- (e) if  $A$  is complex then the above relationships are still true if we use ‘unitary’ in place of ‘orthogonal’ and conjugate transpose in place of transpose. For example, the singular value decomposition of  $A$  is

$$A = QDP^H,$$

where  $Q$  and  $P$  are unitary,  $P^H$  the conjugate transpose of  $P$  and  $D$  is as in (d) above.

## 2.2 Matrix Factorizations

The routines in this section perform matrix factorizations which are required for the solution of systems of linear equations with various special structures. A few routines which perform associated computations are also included.

Other routines for matrix factorizations are to be found in Chapter F03, Chapter F07, Chapter F08 and Chapter F11.

This section also contains a few routines associated with eigenvalue problems (see Chapter F02). (Historical note: this section used to contain many more such routines, but they have now been superseded by routines in Chapter F08.)

## 2.3 Matrix Arithmetic and Manipulation

The intention of routines in this section (sub-chapters F01C and F01Z) is to cater for some of the commonly occurring operations in matrix manipulation, e.g., transposing a matrix or adding part of one matrix to another, and for conversion between different storage formats, e.g., conversion between rectangular band matrix storage and packed band matrix storage. For vector or matrix-vector or matrix-matrix operations refer to Chapter F06.

### 3 Recommendations on Choice and Use of Available Routines

**Note.** Refer to the Users' Note for your implementation to check that a routine is available.

#### 3.1 Matrix Inversion

**Note.** Before using any routine for matrix inversion, consider carefully whether it is really needed.

Although the solution of a set of linear equations  $Ax = b$  can be written as  $x = A^{-1}b$ , the solution should **never** be computed by first inverting  $A$  and then computing  $A^{-1}b$ ; the routines in Chapter F04 or Chapter F07 should **always** be used to solve such sets of equations directly; they are faster in execution, and numerically more stable and accurate. Similar remarks apply to the solution of least-squares problems which again should be solved by using the routines in Chapter F04 rather than by computing a pseudo inverse.

(a) Non-singular square matrices of order  $n$

This chapter describes techniques for inverting a general real matrix  $A$  and matrices which are positive-definite (have all eigenvalues positive) and are either real and symmetric or complex and Hermitian. It is wasteful and uneconomical not to use the appropriate routine when a matrix is known to have one of these special forms. A general routine must be used when the matrix is not known to be positive-definite. In all routines the inverse is computed by solving the linear equations  $Ax_i = e_i$ , for  $i = 1, 2, \dots, n$ , where  $e_i$  is the  $i$ th column of the identity matrix.

Routines are given for calculating the approximate inverse, that is solving the linear equations just once, and also for obtaining the accurate inverse by successive iterative corrections of this first approximation. The latter, of course, are more costly in terms of time and storage, since each correction involves the solution of  $n$  sets of linear equations and since the original  $A$  and its  $LU$  decomposition must be stored together with the first and successively corrected approximations to the inverse. In practice the storage requirements for the 'corrected' inverse routines are about double those of the 'approximate' inverse routines, though the extra computer time is not prohibitive since the same matrix and the same  $LU$  decomposition is used in every linear equation solution.

Despite the extra work of the 'corrected' inverse routines they are superior to the 'approximate' inverse routines. A correction provides a means of estimating the number of accurate figures in the inverse or the number of 'meaningful' figures relating to the degree of uncertainty in the coefficients of the matrix.

The residual matrix  $R = AX - I$ , where  $X$  is a computed inverse of  $A$ , conveys useful information. Firstly  $\|R\|$  is a bound on the relative error in  $X$  and secondly  $\|R\| < \frac{1}{2}$  guarantees the convergence of the iterative process in the 'corrected' inverse routines.

The decision trees for inversion show which routines in Chapter F04 and Chapter F07 should be used for the inversion of other special types of matrices not treated in the chapter.

(b) General real rectangular matrices

For real matrices F08AEF and F01QJF return  $QR$  and  $RQ$  factorizations of  $A$  respectively and F08BEF returns the  $QR$  factorization with column interchanges. The corresponding complex routines are F08ASF, F01RJF and F08BSF respectively. Routines are also provided to form the orthogonal matrices and transform by the orthogonal matrices following the use of the above routines. F01QGF and F01RGF form the  $RQ$  factorization of an upper trapezoidal matrix for the real and complex cases respectively.

F01BLF uses the  $QR$  factorization as described in Section 2.1(ii)(a) and is the only routine that explicitly returns a pseudo inverse. If  $m \geq n$ , then the routine will calculate the pseudo inverse  $Z$  of the matrix  $A$ . If  $m < n$ , then the  $n$  by  $m$  matrix  $A^T$  should be used. The routine will calculate the pseudo inverse  $Z$  of  $A^T$  and the required pseudo inverse will be  $Z^T$ . The routine also attempts to calculate the rank,  $r$ , of the matrix given a tolerance to decide when elements can be regarded as zero. However, should this routine fail due to an incorrect determination of the rank, the singular value decomposition method (described below) should be used.

F02WEF and F02XEF compute the singular value decomposition as described in Section 2 for real and complex matrices respectively. If  $A$  has rank  $r \leq k = \min(m, n)$  then the  $k - r$  smallest singular values will be negligible and the pseudo inverse of  $A$  can be obtained as  $Z = P\Sigma^{-1}Q^T$  as described

in Section 2. If the rank of  $A$  is not known in advance it can be estimated from the singular values (see Section 2.2 of the F04 Chapter Introduction). In the real case with  $m \geq n$ , F02WDF provides details of the  $QR$  factorization or the singular value decomposition depending on whether or not  $A$  is of full rank and for some problems provides an attractive alternative to F02WEF.

### 3.2 Matrix Factorizations

Each of these routines serves a special purpose required for the solution of sets of simultaneous linear equations or the eigenvalue problem. For further details users should consult sections on ‘Recommendations on Choice and Use of Routines’ and ‘Decision Trees’ in the F02 Chapter Introduction and the F04 Chapter Introduction, and individual routine documents.

F01BRF and F01BSF are provided for factorizing general real sparse matrices. For factorizing real symmetric positive-definite sparse matrices, see F11JAF. These routines should be used only when  $A$  is **not** banded and when the total number of non-zero elements is less than 10% of the total number of elements. In all other cases either the band routines or the general routines should be used.

### 3.3 Matrix Arithmetic and Manipulation


The routines in the F01C section are designed for the general handling of  $m$  by  $n$  matrices. Emphasis has been placed on flexibility in the parameter specifications and on avoiding, where possible, the use of internally declared arrays. They are therefore suited for use with large matrices of variable row and column dimensions. Routines are included for the addition and subtraction of sub-matrices of larger matrices, as well as the standard manipulations of full matrices. Those routines involving matrix multiplication may use additional-precision arithmetic for the accumulation of inner products. See also Chapter F06.


The routines in the F01Z section are designed to allow conversion between square storage and the packed storage schemes required by some of the routines in Chapter F02, Chapter F04, Chapter F06, Chapter F07 and Chapter F08.

## 4 Decision Trees

The decision trees show the routines in this chapter and in Chapter F04 that should be used for inverting matrices of various types. Routines marked with an asterisk (\*) only perform part of the computation – see Section 3.1 for further advice.


#### (i) Matrix Inversion:




Tree 1: Inverse of a real  $n$  by  $n$  matrix of full rank

**Note 1:** the inverse of a band matrix  $A$  does not in general have the same shape as  $A$ , and no routines are provided specifically for finding such an inverse. The matrix must either be treated as a full matrix, or the equations  $AX = B$  must be solved, where  $B$  has been initialised to the identity matrix  $I$ . In the latter case, see the decision trees in the F04 Chapter Introduction.

**Note2:** by ‘guaranteed accuracy’ we mean that the accuracy of the inverse is improved by use of the iterative refinement technique using additional precision.

**Tree 2: Inverse of a complex  $n$  by  $n$  matrix of full rank**

Tree 3: Pseudo-inverses



(ii) **Matrix Factorizations:** See the Decision Trees in Section 4 of the F02 Chapter Introduction and Section 4 of the F04 Chapter Introduction.

(iii) **Matrix Arithmetic and Manipulation:** Not appropriate.

## 5 Index

### (i) Inversion

Real $m$ by $n$ Matrix, Pseudo Inverse,	F01BLF
Real Symmetric Positive-definite Matrix, Accurate Inverse,	F01ABF
Real Symmetric Positive-definite Matrix, Approximate Inverse,	F01ADF

### (ii) Matrix Transformations

Complex $m$ by $n$ ( $m \leq n$ ) Matrix, $RQ$ Factorization,	F01RJF
Complex Matrix, Form Unitary Matrix,	F01RKF
Complex Upper Trapezoidal Matrix, $RQ$ Factorization,	F01RGF
Eigenproblem $Ax = \lambda Bx$ , $A, B$ Banded, Reduction to Standard Symmetric Problem,	F01BVF
Tridiagonal matrix, $LU$ Factorization,	F01LEF
Real Almost Block-diagonal Matrix, $LU$ Factorization,	F01LHF
Real Band Symmetric Positive-definite Matrix, $ULDL^T U^T$ Factorization,	F01BUF



Real Band Symmetric Positive-definite Matrix,	
Variable Bandwidth, Cholesky Factorization,	F01MCF
Real $m$ by $n$ ( $m \leq n$ ) Matrix, $RQ$ Factorization,	F01QJF
Real Matrix, Form Orthogonal Matrix,	F01QKF
Real Upper Trapezoidal Matrix, $RQ$ Factorization,	F01QGF
Real Sparse Matrix, Factorization,	F01BRF
Real Sparse Matrix, Factorization, Known Sparsity Pattern,	F01BSF

(iii) **Matrix Arithmetic and Manipulation**

Matrix Addition,	
Real Matrices	F01CTF
Complex Matrices	F01CWF
Matrix Multiplication,	F01CKF
Matrix Storage Conversion	
Packed Triangular $\leftrightarrow$ Square Storage	
Real Matrices	F01ZAF
Complex Matrices	F01ZBF
Packed Band $\leftrightarrow$ Rectangular Storage	
Real Matrices	F01ZCF
Complex Matrices	F01ZDF
Matrix Subtraction,	
Real Matrices	F01CTF
Complex Matrices	F01CWF
Matrix Transpose,	F01CRF

## 6 Routines Withdrawn or Scheduled for Withdrawal

Since Mark 13 the following routines have either been withdrawn or superseded. Those routines indicated by a dagger are still present at Mark 19, but will be omitted at a future date. Advice on replacing calls to these routines is given in the document ‘Advice on Replacement Calls for Withdrawn/Superseded Routines’.

F01AAF	F01ACF	F01AEF	F01AFF	F01AGF	F01AHF
F01AJF	F01AKF	F01ALF	F01AMF	F01ANF	F01APF
F01ATF	F01AUF	F01AVF	F01AWF	F01AXF	F01AYF
F01AZF	F01BCF	F01BDF	F01BEF	F01BNF	F01BPF
F01BQF	F01BTF	F01BWF	F01BXF	F01CAF	F01CBF
F01CDF	F01CEF	F01CFF	F01CGF	F01CHF	F01CLF
F01CMF	F01CNF	F01CPF	F01CQF	F01CSF	F01DAF
F01DBF	F01DCF	F01DDF	F01DEF	F01LBF	F01LZF
F01MAF†	F01NAF	F01QAF	F01QBF	F01QCF	F01QDF
F01QEF	F01QFF	F01RCF	F01RDF	F01REF	F01RFF

## 7 References

- [1] Golub G H and van Loan C F (1996) *Matrix Computations* Johns Hopkins University Press (3rd Edition), Baltimore
- [2] Wilkinson J H and Reinsch C (1971) *Handbook for Automatic Computation II, Linear Algebra* Springer-Verlag
- [3] Wilkinson J H (1965) *The Algebraic Eigenvalue Problem* Oxford University Press, London
- [4] Wilkinson J H (1977) Some recent advances in numerical linear algebra *The State of the Art in Numerical Analysis* (ed D A H Jacobs) Academic Press