CHEM-457: Inorganic Chemistry

Midterm II – April 17th, 2014

This exam is comprised of five questions and is 13 pages in length. Please be sure that you have a complete exam and place your name on each page.

Answer each question to the best of your ability. Partial credit will be awarded where appropriate. You are not permitted to use any supplemental materials other than what is included in this test booklet. Calculators are not needed and their use is not permitted for this exam. **PLEASE DO NOT REMOVE ANY PAGES FROM THIS EXAM EXCEPT FOR THE TWO APENDICIES**.


Write all your answers directly in this test booklet and show all work where necessary.

Good Luck!

1.		_ (12 pts)
2.		_ (20 pts)
3.		_ (10 pts)
4.		_ (24 pts)
5.		_ (34 pts)
Tot	al	

Name:	
1.	(12 Total Points) Please give definitions and an example for each of the following (1 point for each definition & 1 point for each example).
	(a) Arrhenius Acid
	(b) Arrhenius Base
	(c) Brønsted-Lowry Acid
	(d) Brønsted-Lowry Base
	(e) Lewis Acid
	(f) Lewis Base

2. **(20 Total Points)** Consider the following proton transfer reaction and corresponding proton affinities. For this problem, you may assume that entropic factors are negligible such that $\Delta G^{\circ} \approx \Delta H = \Delta PA$.



(a) (3 Points) Is the above reaction spontaneous in aqueous solution? Please explain you answer.

(b) (3 Points) Is the above reaction complete in aqueous solutions? Please explain you answer.

(c) (8 Points) Determine the free energy change for the proton transfer in the gas phase. Is the above reaction spontaneous in the gas phase? Please explain your answer?

Name:	


(d) (3 Points) Which of the four species in the acid-base reaction will be best solvated/stabilized in water? Please explain your answer.

(e) (3 Points) Do you predict the gas phase proton affinity of N,N-dimethylaniline ($Ph-NMe_2$) to be larger or smaller than that of aniline? Please explain your answer.

3. **(10 Total Points)** The sol-gel synthesis of oxide materials is accomplished using hydrated metal ions such as that of titanium.

$$[Ti(OH_2)_6]^{4+} + H_2O \rightarrow [Ti(OH_2)_5(OH)]^{3+} + H_3O^+$$

(a) (4 Points) Label each of the four species in the above reaction as either an acid or base. Please place your labels directly in the boxes below.



(b) (4 Points) Using hard-soft acid base principles, explain whether H₂O or OH¯will be a better base toward Ti⁴+.

(c) (2 Points) Do you expect the equilibrium constant (K) for the above reaction to be greater than or less than 1.0? Please explain your answer.

- 4. **(24 total points)** Nitric oxide, which has the chemical formula NO, is a gaseous molecule that functions as a vasodilator in mammals.
 - (a) (8 Points) Prepare a molecular orbital energy level diagram for NO. Please be sure to label all atomic orbitals and all molecular orbitals. Please be sure to provide sketches that clearly show how the atomic orbitals interact to form MOs. (Note: you may find the table of *Atomic Orbital Ionization Energies* on page 13 of this exam to be useful in constructing your MO diagram).

Name:	
	(b) (2 points) How does your diagram illustrate the difference in electronegativity between N and O?

(c) (4 points) What are the bond order and multiplicity of NO?


Name:	

(d) (4 Points) NO⁺ and NO⁻ are also known species. Calculate the bond orders of these species and compare these values with that of NO. Which of these three species would you predict to have shortest bond? Please explain your answer.

(e) (6 points) The ion NO⁻ can react with H⁺ to form a chemical bond. Do you expect this protonation to occur on the oxygen or nitrogen atom of NO⁻? Do you expect this acid-base adduct to be linear or bent? Please explain your reasoning.

Name:			


5. (34 Total Points) In this problem, let us consider the bonding for methylene (CH₂) using both the 2s and 2p valence orbitals of C and symmetry adapted linear combinations (SALCs) of H 1s orbitals as the basis set. The two H-atom SALCs are illustrated below.



(a) (12 Points) Prepare a molecular orbital energy level diagram for CH₂ in a *linear* geometry. Please be sure to label all atomic orbitals, SALCs and molecular orbitals. Please be sure to provide sketches of the group orbitals and clearly show how they interact with the appropriate orbitals on carbon to form MOs. (Note: you may find the table of *Atomic Orbital Ionization Energies* on page 13 of this exam to be useful in constructing your MO diagram).

(b) (2 Points) What is the bond order of linear methylene? Is this species diamagnetic or paramagnetic?

(c) (12 Points) Prepare a molecular orbital energy level diagram for CH₂ in a **bent** geometry. The appropriate two H-atom SALCs are illustrated below. Please be sure to label all atomic orbitals, SALCs and molecular orbitals. Please be sure to provide sketches of the group orbitals and clearly show how they interact with the appropriate orbitals on carbon to form MOs. (Note: you may find the table of *Atomic Orbital Ionization Energies* on page 13 of this exam to be useful in constructing your MO diagram).



(d) (2 Points) What is the bond order of bent methylene? Is this species diamagnetic or paramagnetic?

Name:	•	

(e) (6 Points) Based on your MO diagrams for linear and bent methylene, which geometry of this species do you expect to be more stable? Please explain your answer.

PERIODIC TABLE OF THE ELEMENTS

232.038	Th a	9	140.12	င္ပ	58
231.036	Pa	01	140.908	P_{r}	59
238.029	٦,	S	144.24	ΡN	8
(237)	Ψ,	02	(145)	$_{ m Pm}$	61
(244)	Pu	0.4	150.36	Sm	62
(243)	Am	08	151.86	Eu	ස
(247)	Cm	08	157.25	Gd	84
(247)	Bk.	07	158.925	Тъ	65
(251)	Cf.	90	162.50	Dу	88
(252)	Es a	8	164.930	Но	67
(257)	Fm	100	167.26	Εŗ	88
(258)	Md	101	168.934	T_{m}	69
(259)	N _E	100	173.04	ΥЪ	70
(260)	ŗ.	102	174.987	Lμ	71

			_			00			ω			2						_	
(223)	Fr	87	132.905	CS	55	85.4678	Rъ	37	39.0983	×	19	22.9898	Na	11	6.941	Li	ω	1.0079	Η
226.025	Ra	88	137.33	Ba	56	87.62	Sr	38	40.08	Ca	20	24.305	Mg	12	9.01218	Be	4		
226.025 227.028	Ac	89	138.906	La	57	88.9059	Υ	39	44.9559	Sc	21							•	
(261)	Unq	104	178.49	$_{ m Hf}$	72	91.224	Zr	40	47.88	Ti	22								
(262)	U_{mp}	105	180.948	Ta	73	92.9064	N	41	50.9415	<	23								
(263)	U_{nh}	106	183.85	W	74	95.94	Mo	42	51.996	$\mathcal{C}^{\mathbf{r}}$	24								
(262)	U_{ns}	107	186.207	Re	75	(98)	Tc	43	54.9380	M_n	25								
(285)	U_{no}	108	190.2	S _o	76	101.07	Ru	44	55.847	Fe	26								
(266)	Une	109	192.22	ŀ	77	102.906	Rh	45	58.9332	င္ပ	27								
			195.08	Ρt	78	106.42	Pd	46	58.69	Z.	28								
			196.967	Au	79	107.868	Αg	47	63.546	Cu	29								
			200.59	Hg	80	112.41	С	48	65.39	Zn	30								
			204.383	Τ1	81	114.82	In	49	69.72	Ga	31	26.9815	Αl	13	10.81	В	51		
			207.2	РЬ	82	118.71	Sn	50	72.59	Ge	32	28.0855	Si	14	12.011	С	o		
			208.980	Βı	83	121.75	SЪ	51	74.9216	As	ಜ	30.9738	P	15	14.0067	Z	7		
			(209)	Ро	8	127.60	Te	52	78.96	Se	34	32.06	S	16	15.9994	0	œ		
			(210)	At	85	126.905 131.29	П	53	79.904	Βr	35	35.453	Ω	17	18.9984 20.179	F	9		
			(222)	Rn	86	131.29	Xe	54	83.80	Κr	36	39.948	Ar	18	20.179	Ne	10	4.00260	He ²

Atomic Orbital Ionization Energies (eV)

Valence Level	Element	1s	2s	2p	3s	3p	4s	4р
1	Н	-13.6						
1	He	-24.5						
2	Li		-5.45					
2	Be		-9.30					
2	В		-14.0	-8.30				
2	С		-19.5	-10.7				
2	N		-25.5	-13.1				
2	0		-32.3	-15.9				
2	F		-46.4	-18.7				
2	Ne		-48.5	-21.5				
3	Na				-5.21			
3	Mg				-7.68			
3	Al				-11.3	-5.95		
3	Si				-15.0	-7.81		
3	Р				-18.7	-10.2		
3	S				-20.7	-11.7		
3	Cl				-25.3	-13.8		
3	Ar				-29.2	-15.9		
4	K						-4.34	
4	Ca						-6.07	
4	Zn						-9.42	
4	Ga						-12.6	-5.95
4	Ge						-15.6	-7.56
4	As						-17.6	-9.05
4	Se						-20.8	-10.8
4	Br						-24.0	-12.5
4	Kr						- 27.5	-14.3

Valence Level	Element	1s	2 s	2p
4	Sc	-4.71	− 5.70	-3.22
4	Ti	- 5.58	-6.07	-3.35
4	٧	-6.32	-6.32	-3.47
4	Cr	- 7.19	-6.57	-3.47
4	Mn	-7.93	-6.82	-3.59
4	Fe	-8.68	-7.07	-3.72
4	Co	-9.42	-7.32	-3.84
4	Ni	-10.0	-7.56	-3.84
4	Cu	-10.7	-7.69	-3.97