


Physical Chemistry

Lecture 7

Special steps; chain reactions;
surface and enzyme kinetics


Photochemical and sensitized steps

◆ Light is sometimes used to activate processes



■ ϕ = quantum yield


◆ Some added materials produce reactive species by reaction



Unimolecular reactions: Lindemann's mechanism


◆ "Simple" reactions are more complex than they seem

◆ Frederick Lindemann proposed intervention of a mediator to produce a highly reactive intermediate in unimolecular reactions



$$\frac{d\text{Products}}{dt} = k_{eff} [A]$$

$$k_{eff} = \frac{k_1 k_3 [M]}{k_3 + k_2 [M]}$$



Variations and refinements of Lindemann's mechanism

◆ Hinshelwood

■ Explicit energy dependence of the rate constant

◆ Rice and Ramsperger, and independently Kassel developed a theory (RRK theory)


■ Explicit account of molecular vibrational state

◆ Marcus

■ Refined RRK theory in a number of ways

◆ Modern version is called RRKM theory

■ Predicts functional dependence of unimolecular reaction rates well



From Engel and Reid, Thermodynamics
Copyright © 2006 Pearson Education, Inc. publishing as Benjamin Cummings

Chain reactions

◆ Many reactions have multiple steps in the mechanism

◆ Chain reactions, once started, continue


- Polymerization
- Some photochemical reactions

◆ Classes of steps

- Initiation - produce reactive species
- Propagation - remove and produce reactive species
- Termination - remove reactive species

Initiation steps


◆ Photochemical steps



Thermal steps





◆ Sensitized steps



Propagation steps


- ◆ Steps that use and create reactive species

- ◆ Examples:



Termination steps

- ◆ Steps that remove reactive species
- ◆ Stable-product formation



- ◆ Wall deactivation




- ◆ Stable-radical formation; scavenging





Vinyl polymers


- ◆ "Simple" chain reaction $n \text{ (monomer)} \rightarrow \text{polymer}$
 $n M \rightarrow M_n$



Vinyl polymerization

- ◆ Chain reaction
- ◆ Generally initiated with some radical
 - Deliberately added
 - Photochemically induced



$$\vdots$$


Mathematics of vinyl polymerization



- ◆ Approximation at steady state
 - Rate of initiation is equal to rate of termination
- ◆ Radical-combination termination

$$v_p = k_p \sqrt{\frac{k_i f}{k_t}} [M][In]^{1/2}$$

- ◆ Other possible termination steps
 - Disproportionation
 - Chain transfer


Reactions at surfaces

- ◆ Very often reaction happens at "special" sites
 - Enzyme action
 - Heterogeneous catalysis
- ◆ Simple surface reaction scheme



Langmuir isotherms

- ◆ Assume equilibrium between gas-phase A and adsorbed A
 - Langmuir isotherm gives relation between gas and surface concentrations



$$[A_{adsorbed}] = [A_{sat}] \frac{bP_A}{1 + bP_A}$$

- Generalize for multiple materials adsorbed, as in a chemical reaction

$$[A_{ads}] = S_0 \frac{b_A P_A}{1 + \sum_j b_j P_j}$$

Langmuir-Hinshelwood kinetics

- ◆ Second-order surface-mediated reaction
- ◆ Rate depends on the partial pressures of A and B

Rate-determining Step



- At low pressure, rate is second-order in the gas pressure
- At high pressures of both reactants, the rate becomes zero-order in pressure

$$v = k_{react} [A_{ads}] [B_{ads}]$$

$$= k_{react} S_0^2 \frac{b_A b_B P_A P_B}{(1 + \sum_j b_j P_j)^2}$$

Eley-Rideal kinetics

- ◆ Second-order surface-mediated transformation

Rate-determining Step

- ◆ One of the reactants comes in from the gas phase (without adsorption)



$$v = k_{react} P_A [B_{ads}]$$

$$= k_{react} S_0 \frac{b_B P_A P_B}{1 + \sum_j b_j P_j}$$

- Always first order with respect to A
- Usually requires a highly reactive gas-phase species such as H atom

Langmuir-Hinshelwood versus Eley-Rideal kinetics

- ◆ Langmuir-Hinshelwood kinetics
 - Both partners of a second-order reaction at the surface
 - Partners diffuse on surface until meeting to react
- ◆ Eley-Rideal kinetics
 - One partner of a second-order reaction held at the surface
 - Second comes directly from the gas phase
 - One or both must be highly reactive



Michaelis-Menten enzyme catalysis

- ◆ Mechanism is similar to surface catalysis

- Form complex
- Complex may



- ◆ Fall apart
- ◆ React



- ◆ Velocity is found assuming fast equilibrium of first two steps

$$v = \frac{k_2}{K_M + [S]} [E]_0 [S]$$


Example of Michaelis-Menten kinetics

- ◆ Hydrolysis of N-glutamyl-L-phenylalanine with α -chymotrypsin

- ◆ J. Chem. Ed., 50, 149 (1973).

- ◆ Lineweaver-Burk plot
- Plot $1/v$ versus $1/[S]$

- ◆ Obtain Michaelis-Menten parameters from slope and intercept of plot



Summary

- ◆ Complex reactions usually described in terms of elementary steps
- ◆ Lindemann's mechanism
 - Modern version is RRKM theory (Rice, Ramsperger, Kassel, and Marcus)
- ◆ Polymerization occurs by a chain reaction
 - Initiation
 - Propagation
 - Termination
- ◆ Surface chemistry
 - Adsorption and desorption steps included
 - Langmuir-Hinshelwood versus Eley-Rideal mechanisms
- ◆ Enzyme kinetics
 - Formation of complex
 - Michaelis-Menten kinetics